Lawrence Livermore National Laboratory (LLNL) is a premier research laboratory that is part of the National Nuclear Security Administration (NNSA) within the U.S. Department of Energy (DOE). As a national security laboratory, LLNL is responsible for ensuring that the nation's nuclear weapons remain safe, secure, and reliable. The Laboratory also meets other pressing national security needs, including countering the proliferation of weapons of mass destruction and strengthening homeland security, and conducts major research in atmospheric, earth, and energy sciences; bioscience and biotechnology; and engineering, basic science, and advanced technology. The Laboratory serves as a scientific resource to the U.S. government and a partner to industry and academia.

Since its inception in 1952 until October 1, 2007, LLNL was managed by the University of California. In May 2007, DOE selected Lawrence Livermore National Security, LLC (LLNS), to manage the Laboratory under a seven-year contract that began on October 1, 2007.

LLNL operations release a variety of constituents into the environment via atmospheric, surface water, and groundwater pathways. Some of the constituents, such as particles from diesel engines, are common at many types of facilities while others, such as radionuclides, are unique to facilities like LLNL. All releases are highly regulated and carefully monitored.

Safe, secure, and efficient operations that provide a safe, clean environment for employees and neighboring communities are a necessary part of the Laboratory's research and development programs and underpin their success. Experts in environment, safety and health (ES&H) support all Laboratory activities. LLNL's radiological control program ensures that radiological exposures and releases are reduced to as low as reasonably achievable to protect the health and safety of its employees, contractors, the public, and the environment.

LLNL is committed to enhancing its environmental stewardship and reducing any impacts its operations may have on the environment. The Laboratory encourages the public to participate in matters related to the Laboratory's environmental impact on the community by soliciting citizens' input on matters of significant public interest and through various communications. The Laboratory also provides public access to information on its ES&H activities.

LLNL consists of two sites—an urban site in Livermore, California, referred to as the "Livermore site," which occupies 1.3 square miles; and a rural Experimental Test Site, referred to as "Site 300," near Tracy, California, which occupies 10.9 square miles. In 2007 the Laboratory had a staff of more than 8000.

Purpose and Scope of the Environmental Report

The purposes of the *Environmental Report 2007* are to record LLNL's compliance with environmental standards and requirements, describe LLNL's environmental protection and remediation programs, and present the results of environmental monitoring. Specifically, the

report discusses LLNL's Environmental Management System; describes significant accomplishments in pollution prevention; presents the results of air, water, vegetation, and foodstuff monitoring; reports radiological doses from LLNL operations; summarizes LLNL's activities involving special status wildlife, plants, and habitats; and describes the progress LLNL has made in remediating groundwater contamination.

Environmental monitoring at LLNL, including analysis of samples and data, is conducted according to LLNL's Environmental Protection Department Quality Assurance Management Plan, which is based on DOE Order 414.1C, Quality Assurance.

This report is prepared for DOE by LLNL's Environmental Protection Department. Submittal of the report satisfies requirements under DOE Order 231.1A, Environmental Safety and Health Reporting, and DOE Order 5400.5, Radiation Protection of the Public and Environment. The report is distributed in electronic form and is available to the public at https://saer.llnl.gov/, the website for the LLNL annual environmental report. Previous LLNL annual environmental reports beginning in 1994 are also on the website.

Regulatory Permitting and Compliance

LLNL undertakes substantial activities to comply with many federal, state, and local environmental laws. The major permitting and regulatory activities that LLNL conducts are required by the Clean Air Act; the Clean Water Act and related state programs; the Emergency Planning and Community Right-to-Know Act, the Resource Conservation and Recovery Act and state and local hazardous waste regulations; the National Environmental Policy Act and the California Environmental Quality Act; the Endangered Species Act; the National Historic Preservation Act; the Antiquities Act; and the Comprehensive Environmental Response, Compensation and Liability Act.

Environmental Management System

LLNL established its Environmental Management System (EMS) to meet the requirements of the International Organization for Standardization (ISO) 14001:1996 in June 2004. In June 2006, LLNL upgraded its EMS to meet the requirements of ISO 14001:2004. In late 2006, the EMS was extended to the directorate level.

To further implement LLNL's EMS, a Multi-Directorate Consortium was formed to provide a forum for identifying common environmental issues. The Consortium was a venue for presentations on shared chemical usage, energy topics, and office paper use and reduction. During 2007, six directorates completed one or more directorate environmental management plans (EMPs) and the EMPs developed to address Lab-wide environmental aspects during 2006 are still in progress or contain an ongoing component.

Pollution Prevention

A strong Pollution Prevention (P2) Program is an essential element of LLNL's EMS. The P2 Team is responsible for P2 program stewardship and maintenance, waste stream analysis, waste generation reporting, and coordination of institutional P2 programs and activities.

Two LLNL projects were selected by NNSA/Headquarters to receive 2007 Environmental Stewardship awards. The first award was for the Space Action Team's "Assets for Value" process, which allows the reuse/salvage value of materials to be included as an offset to bids for decontamination and demolition. The second award was to Fleet Management for the construction and operation of an E85 fuel dispensing station.

LLNL also conducted activities to promote employee awareness of P2, including the annual Earth Expo held in April to coincide with Earth Day, articles in the LLNL newspaper, and training for procurement staff.

Air Monitoring

LLNL operations involving radioactive materials had minimal impact on ambient air during 2007. Estimated nonradioactive emissions are small compared to local air district emission criteria.

Releases of radioactivity to the environment from LLNL operations occur through stacks and from diffuse area sources. In 2007, radioactivity released to the atmosphere was monitored at six facilities on the Livermore site and one at Site 300. In 2007, 0.57 TBq (15.4 Ci) of tritium was released from the Tritium Facility, and 1.7 GBq of tritium (46 mCi) was released from the Decontamination and Waste Treatment Facility. The Contained Firing Facility at Site 300 had 2.1 kBq (57 nCi) of depleted uranium released in particulate form in 2007. None of the other facilities monitored for gross alpha and gross beta radioactivity had emissions in 2007.

The magnitude of nonradiological releases (e.g., reactive organic gases/precursor organic compounds, nitrogen oxides, carbon monoxide, particulate matter, sulfur oxides) is estimated based on specifications of equipment and hours of operation. Estimated releases in 2007 for the Livermore site were of similar levels to those in 2006; estimated releases at Site 300 were higher in 2007 than in 2006 due primarily to the required periodic preventative maintenance of emergency stand-by diesel generators. Nonradiological releases from LLNL continue to be a very small fraction of releases from all sources in the Bay Area or San Joaquin County.

In addition to air effluent monitoring, LLNL samples ambient air for tritium, radioactive particles, and beryllium. Some samplers are situated specifically to monitor areas of known contamination; some monitor potential exposure to the public; and others, distant from the two LLNL sites, monitor the natural background. In 2007, ambient air monitoring data confirmed estimated releases from monitored stacks and were used to determine source terms for resuspended plutonium-contaminated soil and tritium diffusing from area sources at the Livermore site and

resuspended uranium-contaminated soil at Site 300. In 2007, radionuclide particulate, tritium, and beryllium concentrations in air at the Livermore site and in the Livermore Valley were well below the levels that would cause concern for the environment or public health.

Water Monitoring

Monitoring of various categories of water is carried out to determine whether any radioactive or nonradioactive constituents released by LLNL might have a negative impact on public health and the environment. Data indicate LLNL has good control of its discharges to the sanitary sewer, and current discharges to the surface water and groundwater do not have any significant environmental impact.

Permits, including one for discharging treated groundwater from the Livermore site Ground Water Project, regulate discharges to the City of Livermore sanitary sewer system. During 2007, no discharges to the sanitary sewer exceeded any effluent limits for radioactive materials, and all the values were less than 2% of the allowed limits. For nonradioactive materials, all constituents were less than 20% of the allowed limits. All discharges to the Site 300 sewage evaporation and percolation ponds were within permitted limits, and groundwater monitoring showed no measurable impacts.

Storm water is sampled for constituents such as radioactivity, metals, oxygen, dioxins, polychlorinated biphenyls (PCBs), and nitrate both upstream and downstream from both the Livermore site and Site 300. In 2007, no acute or chronic toxicity was seen in runoff, and data showed that the quality of Livermore site storm water effluent was similar to that entering the site (influent). At Site 300 in 2007, data continued to show that concentrations remained below levels of environmental concern.

Extensive monitoring of groundwater occurs at and near the Livermore site and Site 300. Groundwater from wells downgradient from the Livermore site is analyzed for pesticides, herbicides, radioactivity, nitrates, and hexavalent chromium. To detect any off-site contamination quickly, the well water is sampled in the uppermost water-bearing layers. Near Site 300, monitored constituents in off-site groundwater include explosives residue, nitrate, perchlorate, metals, volatile and semivolatile organic compounds, tritium, uranium, and other (gross alpha and beta) radioactivity. With the exception of volatile organic compounds (VOCs) in wells monitored for CERCLA compliance, the constituents of all off-site samples collected at both the Livermore site and Site 300 in 2007 were below allowable limits for drinking water.

Surface waters and drinking water are analyzed for tritium and gross alpha and gross beta radioactivity. In the Livermore Valley, the maximum tritium activity was less than 1% of the drinking water standard, and the maximum gross alpha and gross beta measurements were less than 35% of their respective drinking water standards. For Lake Haussmann (formerly called the Drainage Retention Basin) on the Livermore site, levels of gross alpha, gross beta, tritium, metals, and pesticides were below discharge limits, and organics and PCBs were below detection limits. pH exceedance of the discharge limits correspond to seasonal algal blooms. Aquatic

bioassays for acute and chronic toxicity showed no effects in water discharged from Lake Haussmann. At Site 300, discharges from maintenance and the operation of drinking water system and mechanical equipment resulted in discharges that were below permitted limits.

Terrestrial Radiological Monitoring

The impact of LLNL operations on surface soil, sediment, and vadose zone soils in 2007 was insignificant. Soils and sediments are analyzed for plutonium, gamma-emitting radionuclides, tritium, total and soluble metals, and PCBs as appropriate. Plutonium concentrations at the Livermore Water Resources Division (formerly called the Livermore Water Reclamation Plant) continued to be high relative to other sampled locations, but even this concentration was only 1.5% of the screening level for cleanup recommended by the National Council on Radiation Protection (NCRP). At Site 300, soils are analyzed for gamma-emitting radionuclides and beryllium. In 2007, uranium-238 concentrations in soils at Site 300 were below NCRP-recommended screening levels. Beryllium concentrations were within the ranges reported since sampling began in 1991.

Vegetation and Livermore Valley wine were sampled for tritium. In 2007, the median concentrations in all off-site vegetation samples were below the lower limit of detection of the analytical method. The highest concentration of tritium in Livermore Valley wines sampled in 2007 was 0.3% of the drinking water standard.

LLNL's extensive network of thermoluminescent dosimeters and real-time sensors measure the natural terrestrial and cosmogenic background; in 2007, as in recent years, no impact from LLNL operations was detected.

Biota

Through monitoring and compliance activities in 2007, LLNL avoided most impacts to special status species and enhanced some habitats. LLNL studies, preserves, and tries to improve the habitat of five species at Site 300 that are covered by the federal or California Endangered Species Acts—California tiger salamander (*Ambystoma californiense*), California red-legged frog (*Rana aurora draytonii*), Alameda whipsnake (*Masticophus lateralis euryxanthus*), valley elderberry longhorn beetle (*Desmocerus californicus dimorphus*), and the large-flowered fiddleneck (*Amsinckia grandiflora*)—as well as species that are rare and otherwise of special interest. At Site 300, LLNL monitors populations of birds and rare species of plants and also continues restoration activities for the four rare plant species known to occur at Site 300—the large-flowered fiddleneck, the big tarplant (*Blepharizonia plumosa*, also known as *Blepharizonia plumosa* subsp *plumosa*), the diamond-petaled poppy (*Eschscholzia rhombipetala*), and the round-leaved filaree (*Erodium macrophyllum*).

LLNL took several actions to control invasive species in 2007. Measures taken at the Livermore site to control bullfrogs, which are a significant threat to California red-legged frogs, included

dispatching adults, removing egg masses, and allowing part of Arroyo Las Positas to dry out in October 2007. Site 300's invasive species control efforts have been focused largely on dispatching feral pigs. In 2007, six feral pigs were dispatched.

The 2007 radiological doses calculated for biota at the Livermore site or Site 300 were far below screening limits set by DOE, even though highly conservative assumptions maximized the potential effect of LLNL operations on biota.

Radiological Dose

Annual radiological doses at the Livermore site and Site 300 in 2007 were found to be well below the applicable standards for radiation protection of the public. Dose calculated to the site-wide maximally exposed individual (SW-MEI) for 2007 was 0.031 μ Sv (0.0031 mrem) for the Livermore site and 0.035 μ Sv (0.0035 mrem) at Site 300. These doses are well below the federal National Emissions Standards for Hazardous Air Pollutants of 100 μ Sv (10 mrem) and are significantly less than the doses from natural background radiation (see **Section 7.3.4**). Four sources of tritium at LLNL contributed nearly 100% of the dose received by the SW-MEI. There were no unplanned releases of radionuclides to the atmosphere at the Livermore site or at Site 300.

Groundwater Remediation

Groundwater at both the Livermore site and Site 300 is contaminated from historical operations; the contamination, for the most part, is confined to each site. Groundwater at both sites is undergoing cleanup under the Comprehensive Environmental Response, Compensation and Liability Act (CERCLA). Remediation activities removed contaminants from groundwater and soil vapor at both sites, and documentation and investigations continue to meet regulatory milestones.

At the Livermore site, contaminants include VOCs, fuel hydrocarbons, metals, and tritium, but only the VOCs in groundwater and saturated and unsaturated soils need remediation. VOCs are the main contaminant found at the nine Site 300 operable units (OUs). In addition, nitrate, perchlorate, tritium, high explosives, depleted uranium, organosilicate oil, and metals are found at one or more of the OUs.

In 2007, concentrations continued to decrease in most of the Livermore site VOC plumes due to active remediation and the removal of more than 318 kg of VOCs from both groundwater and soil vapor. VOC concentrations on the western margin of the site continued their decline, indicating effective hydraulic control of the boundary plumes. In the interior of the site, remediation activities, including soil vapor extraction, dual extraction, and groundwater extraction, have resulted in declines of VOC concentrations in source areas.

In 2007 at Site 300, perchlorate, nitrate, the high explosive RDX, and organosilicate oil were removed from groundwater in addition to about 62 kg of VOCs. Each Site 300 OU has a different profile of contaminants, but overall, groundwater and soil vapor extraction and natural attenuation continue to reduce the mass of contaminants in the subsurface. Cleanup remedies have been fully implemented and are operational at seven of the nine OUs at Site 300; cleanup remedies for one of the remaining units has been submitted and the Remedial Investigation/Feasibility Study for the other OU is scheduled for 2008.

Conclusion

The combination of surveillance and effluent monitoring, source characterization, and dose assessment showed that the radiological dose to the hypothetical, most-exposed member of the public caused by LLNL operations in 2007 was substantially less than the dose from natural background. Potential dose to biota was well below DOE screening limits. LLNL demonstrated good compliance with permit conditions for releases to air and to water. Analytical results and evaluations of air and various waters potentially impacted by LLNL operations showed minimal contributions from LLNL operations. Remediation efforts at both the Livermore site and Site 300 further reduced concentrations of contaminants of concern in groundwater and soil vapor.